AD-A114 133

TEXAS UNIV AT AUSTIN CENTER FOR CYBERNETIC STUDIES

HANDLING ARTIFICIAL VARIABLES IN A SIMPLEX METHOD: A COMMENTARY--ETC(U)

NOV 81 A CHARNES? W W COOPER

UNCLASSIFIED

CC5-415

END

RMD

RMD

RMD

S82

OTIE

AD A 1 14 133

## CENTER FOR CYBERNETIC STUDIES

The University of Texas Austin, Texas 78712

DTIC ELECTE WAY 5 1982

M STATESCAT A

for public releases



82 05 03 09**3** 

(2)

Research Report CCS 415

HANDLING ARTIFICIAL VARIABLES IN A SIMPLEX METHOD: A COMMENTARY

bу

A. Charnes W.W. Cooper

November 1981



This research was partly supported by Office of Naval Research Contract N00014-75-C-0569 with the Center for Cybernetic Studies, The University of Texas at Austin. Reproduction in whole or in part is permitted for any purpose of the United States Government.

CENTER FOR CYBERNETIC STUDIES

A. Charnes, Director
Business-Economics Building, 203E
The University of Texas at Austin
Austin, Texas 78712
(512) 471-1821



## Handling Artificial Variables in A Simplex Method: A Commentary by A. Charnes and W.W. Cooper

Charnes' non-Archimedean Simplex method is unfortunately misrepresented in the paper by P. Jolly "Handling Artificial Variables in the Simplex Method", <u>JIOS</u>, v. 2 (1981), No. 3, pp. 241-248, possibly because he refers to books by Gass [1], 1969, and Hadley [2], 1973, rather than to a correct description as in Charnes and Cooper [3], 1961.

The latter, available as a text, describes the treatment in detail, see also Charnes, Cooper and Henderson [4], 1953, although the general format has been available since the original publication by Charnes in Econometrica [5], 1952. Thus, referring to the former, the use of the non-Archimedean transcendental M can be accomplished by calculation solely in the base field by use of two rows for the  $(z_j - c_j)$ , one of which contains the coefficients of M. There is thus no difficulty with hand computation since each row transfers from one basis to the next as if there were simply one additional row in the tableau.

One could employ this procedure just as well in the machine computation case at the cost of only slightly increasing the machine operations in examining  $(z_j - c_j)$ -values.


Contrary to Jolly's statements on computational experience attributed to Hadley, the use of "large"numbers for M does not pose substantial difficulties for machine computation, when this course, rather than the a priori exact course above is followed. For sufficiently large numbers one obtains the same optimal solution as if the exact course were followed.

Very often, also, one can identify m -1 structural vectors linearly independent of the right hand side vector Po. One can then start with these

in a basis of vectors in which Po is the only artificial one. Any decent LP code today would take this basis and immediately invert it, thus starting the simplex procedure with only one artificial vector to be removed.

## REFERENCES

- 1. S.I. Gass, Linear Programming, Methods and Applications, McGraw-Hill Kogakusha Ltd, 1969.
- 2. G. Hadley, Linear Programming, Addison-Wesley, 1973.
- 3. A Charnes and W.W. Cooper, Management Models and Industrial Applications of Linear Programming, vols. I and II, J. Wiley and Sons, 1961.
- 4. A Charnes, W.W. Cooper, A. Henderson, An Introduction to Linear Programming, J. Wiley and Sons, 1953.
- 5. A. Charnes, "Optimality and Degeneracy in Linear Programming", Econometrica, v. 20, No. 2, pp. 160-170, April, 1952.



According For



Unclassified
SECURITY CLASSIFICATION OF THIS PAGE (When Date Entered)

REPORT DOCUMENTATION PAGE		READ DISTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER	2. GOVT ACCESSION NO.	
CCS 415	AD-A1142	92
4. TITLE (and Subtitle)		5. TYPE OF REPORT & PERIOD COVERED
Handling Artificial Variables in a Simplex		
Method: A Commentary		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(a)		8. CONTRACT OR GRANT NUMBER(s)
AUTHOR(s)		
A. Charnes, W.W. Cooper		N00014-75-C-0569
9. PERFORMING ORGANIZATION NAME AND ADDRESS		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT HUMBERS
Center for Cybernetic Studies, UT Austin		
Austin, TX 78712		
11. CONTROLLING OFFICE NAME AND ADDRESS		12. REPORT DATE
Office of Naval Research (Code 434) Washington, DC		November 1981
		3
14. MONITORING AGENCY NAME & ADDRESS(If different from Controlling Office)		18. SECURITY CLASS. (of this report)
		Unclassified
		18a. DECLASSIFICATION/DOWNGRADING
16. DISTRIBUTION STATEMENT (of this Report)		
This document has been approved for public release and sale; its		
distribution is unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)		
Charnes' non-Archimedean Simplex Method Artificial Variables		
non-Archimedean Transcendentals		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number)		
Charnes' non-Archimedean simplex method and its computation in terms		
of Archimedean quantities is developed to correct erroneous statements		
in the research literature centering around methods of computation involving artificial variables.		
Λ.		
	,	

